

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

VIII. An Account of an Experiment, concerning the Angle requir d to suspend a Drop of Oyl of Oranges, at certain Stations, between two Glass Planes, placed in the form of a Wedge. By Mr. Fr. Hauskbee, F. R. S.

Procured two Glass Planes that measured a Radius of twenty Inches each; their breadth was about 3 Inches: That which I used for the lower Plane, was placed with its Surface parallel with the Center of its Axis, and parallel with the Horizon. Thus (the Planes being very clean) they were rubb'd with a clean Linnen Cloth dipt in Oyl of Oranges: Then a drop or two of the fame Oyl being let fall on the lower Plane near the Axis, the other Plane was laid on it; which so soon as it touch'd the Oyl, the Oyl spread itself considerably between both Then the upper Plane being raised a their Surfaces. little at the same end by a Screw, the Oyl immediately attracted itself into a Body, forming a Globule contiguous to both Surfaces, and began to move forwards toward the touching ends. When it had arriv'd two Inches from the Axis, an Elevation of 15 Minutes at the touching ends stopt its progress, and it remain'd there without Motion any way. The Planes being let fall again, the drop mov'd forward till it came to four Inches from the Center; then an Elevation of 25 Minutes was requir'd to give it a fixt Station. At 6 Inches it requir'd an Angle of 35 Minutes; at 8, of 45 Minutes; at 10, a Degree. At 12 Inches from the Axis, the Elevation was 1 Degree 45 Minutes; and so on, at the several Stations, as they stand in the following Table. This, after abun-

(474)

abundance of tryals, I take to be the most correct, tho' the others succeeded very little different from the same, It is to be observ'd, that when the Globule, or Drop. had arriv'd to near 17 Inches on the Planes from their Axis, it would become of an Oval form; and as it as cended higher, so would its Figure become more and more oblong; and unless the Drop was small, upon fuch an Elevation of the Planes as was requir'd at such a progress of the Drop, it would be parted, some of it descending, and the rest of it running up to the top at once: But upon a Drop that separated thereabouts, I found the remaining part of it at 18 Inches, would bear an Angle of Elevation equal to 22 Degrees to ballance the weight of it. Higher than that I could not obferve. The Planes were separated at their Axis about of an Inch. I found but little difference between small and larger Drops of the Oyl, in relation to the Experiment. The Angles were measured by a Quadrant mark'd on Paper of near 20 Inches Radius, divided into Degrees and Quarters.

Distance in Inches from the Axis.			Ang Elev D.	gle of vation <i>M</i> .
2	-	 	0 -	~15
4	P		0 -	- 25
6	-		0	-35
8		-	0 -	-45
10	-		Ĭ	-00
12	-	-	1	-45
14		-	2 -	-45
15	-		4 -	-00
16	-	خسيسي	6 -	-00
17	-		10-	-00
18	-	-	22-	-00